

TARGET 02 ANSWER KEY

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	2	4	2	1	3	3	2	1	1	3	2	2	4	2	1	1	2	3	1	1
Que.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans.	3	4	3	1	4	4	2	1	3	4	3	2	3	2	2	1	3	2	4	2
Que.	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans.	1	2	2	2	2	3	2	3	3	2	1	1	2	4	3	1	1	2	1	1
Que.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
Ans.	2	2	4	4	4	1	2	4	2	3	3	1	3	3	3	4	3	4	2	2
Que.	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
Ans.	3	3	1	2	1	2	4	2	2	3	4	1	2	2	1	4	3	3	1	3
Que.	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
Ans.	3	3	2	2	1	1	1	1	1	3	2	2	3	2	3	4	4	4	2	3
Que.	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
Ans.	4	2	4	1	1	1	2	3	2	3	1	3	4	2	2	4	3	1	1	4
Que.	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
Ans.	1	3	4	2	3	4	2	1	3	2	3	4	2	4	3	2	3	2	2	3
Que.	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
Ans.	2	4	1	2	2	1	3	4	1	1	4	2	3	3	4	4	2	1	1	1

2. $(2\hat{i} + 3\hat{j} + 2\hat{k}) \cdot (4\hat{i} - 4\hat{j} + \alpha\hat{k}) = 0$
 $8 - 12 + 2\alpha = 0 \Rightarrow \alpha = 2$

3. $R = 2a \cos\theta/2 = 2 \times 6 \times \cos 30^\circ = 6\sqrt{3}$

4. $R = \frac{u^2 \sin 2\theta}{g}$ so that $R_{\max} = \frac{u^2}{g} = 16 \text{ km}$

$$H_{\max} = \frac{u^2 \sin^2 \theta}{2g} = 2 \text{ km}$$

5. $f_L = \mu_s N = 0.54 \times 2 \times 10 = 10.8 \text{ N}$
 $F_{\text{applied}} = 2.8 \text{ N}$
 $\therefore F_{\text{applied}} < f_L$
 $\therefore \text{friction force} = 2.8 \text{ N}$

6. $T - 100g = 100a \quad \dots(i)$

$$T - 60g = 60 \left(\frac{5g}{4} - a \right) \quad \dots(ii)$$

from (i) and (ii) $T = 1218 \text{ N}$

7. $\Delta KE = \frac{1}{2} m [5gr - gr] = 2mgr = 20 \text{ J}$

8. $W = \Delta K$

$$25 = \frac{1}{2} mv^2 = \frac{1}{2} \times 2 \times v^2$$

$$v = 5 \text{ m/s}$$

9. $5u + 0 = (5 + 2.5)v$


$$v = \frac{2}{3}u$$

$$K = \frac{1}{2} \times (5 + 2.5)v^2 = 5 \Rightarrow u = \sqrt{3} \text{ m/s}$$

$$K_1 = \frac{1}{2} \times 5 \times 3 = 7.5 \text{ J}$$

10. under mutual attraction, the centre of mass remains at rest.

15. $T \sin\theta = F_0$
 $T \cos\theta = mg$



$$\tan\theta = \frac{F_0}{mg}$$


$$F_0 = mg \tan\theta \quad \dots(i)$$

In medium

$$\frac{F_0}{K} = mg \left(1 - \frac{\sigma}{\rho}\right) \tan\theta \quad \dots(ii)$$

from (i) & (ii)

$$K = \frac{\rho}{\rho - \sigma}$$



$$\int dF = \int_d^{d+L} \frac{1}{4\pi\epsilon_0} \left(\frac{Q}{L} dx\right) \frac{q}{x^2}$$

$$\Rightarrow F = \frac{1}{4\pi\epsilon_0} \frac{qQ}{L} \left[\frac{1}{-x} \right]_d^{d+L}$$

$$F = \frac{1}{4\pi\epsilon_0} \frac{Qq}{L} \left[-\frac{1}{d+L} + \frac{1}{d} \right]$$


$$= \frac{1}{4\pi\epsilon_0} \frac{qQ}{L} \left[\frac{-d + d + L}{d(d+L)} \right]$$

$$F = \frac{1}{4\pi\epsilon_0} \frac{qQ}{d(d+L)}$$

17. $E = E_{\max} \Rightarrow x = \frac{R}{\sqrt{2}}$ as $\frac{dE}{dx} = 0$

$$E_{\text{axis}} = \frac{1}{4\mu\epsilon_0} \frac{Qx}{(x^2 + R^2)^{3/2}}$$

$$\therefore E_{\max} = \frac{2}{3\sqrt{3}} \left(\frac{1}{4\pi\epsilon_0} \frac{Q}{R^2} \right)$$



At $t = 0$

$$I = \frac{V}{R_{\text{net}}} = \frac{10}{6 + 8} = \frac{10}{14} = \frac{5}{7} \text{ A}$$


33. $11 \times 2\pi \sqrt{\frac{l_s}{g}} = 9 \times 2\pi \sqrt{\frac{l_\ell}{g}}$

$$121 l_s = 81 l_\ell$$

$$\boxed{\frac{l_s}{l_\ell} = \frac{81}{121}}$$

11 oscillation

9 oscillation




34. $A = 6$
at $t = 0$

Particle is at $-\frac{A}{2}$ & moving towards negative extreme.

$$\text{So, } (x - 3) = 6 \sin(4\pi t + 7\pi/6)$$

$$x = 3 + 6 \sin(4\pi t + 7\pi/6)$$



35. $f_c = \frac{(2m+1)V}{4L_c}$

$$f_o = \frac{(m+1)V}{2L_o}$$

For $m = 1$ (1st overtone)

$$f_c = f_o$$

$$\frac{3V}{4L_c} = \frac{2V}{2L_o} \Rightarrow$$

$$\boxed{\frac{L_c}{L_o} = \frac{3}{4}}$$

72. Rate = $k_5 [\text{COCl}]^1 [\text{Cl}_2]^1$ (show step)

$$\frac{k_3}{k_4} = \frac{[\text{COCl}]}{[\text{Cl}][\text{CO}]} \Rightarrow [\text{COCl}] = \frac{k_3}{k_4} [\text{Cl}][\text{CO}]$$

$$\frac{k_1}{k_2} = \frac{[\text{Cl}]^2}{[\text{Cl}_2]} \Rightarrow [\text{Cl}] = \left(\frac{k_1}{k_2} \right)^{1/2} [\text{Cl}_2]^{1/2}$$

$$\text{Rate} = k_5 \times \frac{k_3}{k_4} \times [\text{Cl}] [\text{CO}] [\text{Cl}_2]^1$$

$$\text{Rate} = k_5 \times \left(\frac{k_1}{k_2} \right)^{1/2} [\text{Cl}_2]^{1/2} [\text{CO}] [\text{Cl}_2]^1$$

$$= k_5 \times \frac{k_3}{k_4} \times \left(\frac{k_1}{k_2} \right)^{1/2} \times [\text{CO}]^1 [\text{Cl}_2]^{3/2}$$

73. $A \rightarrow nB$

$A_0 \quad 0$

$A_0 - x \quad nx$

at Intersection $A_0 - x = nx$

$$x = \frac{A_0}{(n+1)}$$

$$\text{Conc. of B} = nx = \frac{nA_0}{(n+1)}$$

74. $\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$

Initial	1	1	0
	$1-\alpha$	$1-3\alpha$	2α

$$1 - 3\alpha = x$$

$$\alpha = \frac{1-x}{3}$$

$$\text{mole of NH}_3 = \frac{2\alpha}{3} = \frac{2(1-x)}{3}$$

75. Sb_2S_3 is negative Sol, So the ion will coagulate the solⁿ, higher the charge on cation higher is coagulative power

92. NCERT-XI Eng. Page No. 86

97. NCERT Pg.# 31,34,37,39

100. NCERT Pg.# 7

101. NCERT Pg.# 38,39

104. NCERT Page No. # 170

109. NCERT-XI, Page 128, Para-2 line-3

110. NCERT-XI, Page 132, Para-4 line-5,6

111. NCERT-XI, Page 133, Para-1

112. NCERT-XI, Page 148, Para-2 line-11,12,13

113. NCERT-XI, Page 128, figure-9.2

114. NCERT-XI, Page 169, figure-10.3

116. NCERT Pg.#169

117. NCERT Pg.#204

118. NCERT Pg.#180

119. NCERT Pg#179

121. NCERT Pg.#247

122. NCERT Pg.#221

123. NCERT Pg.#221

127. NCERT XI Pg. # 262

133. NCERT Pg. # 335(22.2.7)

134. NCERT Pg. # 335(22.2.7)

139. NCERT XII, Pg.# 21

142. NCERT XII, Pg.# 25

144. NCERT Pg. # 61

145. NCERT Pg. # 60

148. 11th Biology Pg. # 147

149. 12th NCERT Pg. # 71

150. 12th NCERT Pg. # 70

156. 12th NCERT Pg. # 100

157. 12th NCERT Pg. # 199

160. 12th NCERT Pg. # 174

161. 12th NCERT Pg. # 187

162. 12th NCERT Pg. # 212 & 213

164. NCERT Pg. # 141

165. NCERT Pg. # 131